PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 H 1.008																	2 He 4.002
3 Li 6.941	4 Be 9.012											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18
11 Na 22.99	12 Mg 24.30											13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.06	17 Cl 35.45	18 Ar 39.95
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.00	25 Mn 54.38	26 Fe 55.85	27 Co 58.93	28 Ni 58.70	29 Cu 63.55	30 Zn 65.39	31 Ga 69.72	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc 98	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 127.9	54 Xe 131.3
55 Cs 132.9	56 Ba 137.3	57 La 138.9	72 Hf 178.5	73 Ta 181.0	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po 209	85 At 210	86 Rn 222
87 Fr 223	88 Ra 226.0	89 Ac 227.0	104 Rf	105 Db	106 Sg	107 Bh	108 Hs	109 Mt	110 Uun								

58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm 145	62 Sm 150.4	63 Eu 152.0	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0	71 Lu 175.0
90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np 237.0	94 Pu 244	95 Am 243	96 Cm 247	97 Bk 247	98 Cf 251	99 Es 252	100 Fm 257	101 Md 258	102 No 259	103 Lr 260

Representative proton nmr chemical shifts





IR CORRELATION CHART

Group		Frequency Range (cm^{-1})	Intensity[a]
A. Alkyl			
C—H (stretching)		2853–2962	(m–s)
Isopropyl, —$CH(CH_3)_2$		1380–1385	(s)
	and	1365–1370	(s)
tert-Butyl, —$C(CH_3)_3$		1385–1395	(m)
	and	~1365	(s)
B. Alkenyl			
C—H (stretching)		3010–3095	(m)
C=C (stretching)		1620–1680	(v)
R—CH=CH_2	(out-of-plane C—H bendings)	985–1000	(s)
	and	905–920	(s)
$R_2C=CH_2$		880–900	(s)
cis-RCH=CHR		675–730	(s)
trans-RCH=CHR		960–975	(s)
C. Alkynyl			
≡C—H (stretching)		~3300	(s)
C≡C (stretching)		2100–2260	(v)
D. Aromatic			
Ar—H (stretching)		~3030	(v)
Aromatic substitution type (C—H out-of-plane bendings)			
Monosubstituted		690–710	(very s)
o-Disubstituted	and	730–770	(very s)
m-Disubstituted		735–770	(s)
		680–725	(s)
	and	750–810	(very s)
p-Disubstituted		800–860	(very s)
E. Alcohols, Phenols, and Carboxylic Acids			
O—H (stretching)			
Alcohols, phenols (dilute solutions)		3590–3650	(sharp, v)
Alcohols, phenols (hydrogen bonded)		3200–3550	(broad, s)
Carboxylic acids (hydrogen bonded)		2500–3000	(broad, v)
F. Aldehydes, Ketones, Esters, and Carboxylic Acids			
C=O (stretching)		1630–1780	(s)
Aldehydes		1690–1740	(s)
Ketones		1680–1750	(s)
Esters		1735–1750	(s)
Carboxylic acids		1710–1780	(s)
Amides		1630–1690	(s)
G. Amines			
N—H		3300–3500	(m)
H. Nitriles			
C≡N		2220–2260	(m)

[a]Abbreviations: s = strong, m = medium, w = weak, v = variable, ~ = approximately.

pK_a TABLE

Acid	pK_a
HI	-9
HBr	-8
HCl	-7
H_2SO_4	-5.2
H_3O^+	-1.74
H_3PO_4	2.1
CH_3CO_2H	4.76
H_2CO_3	6.36
NH_4^+	9.24
HCO_3^-	10.33
H_2O	15.7
CH_3CH_2OH	15.9
$CH_3C{=}OMe$	20
$HC{\equiv}CH$	25
H_2	35
NH_3	38
$CH_2{=}CH_2$	44
CH_3CH_3	51